

ANSWERS

1

Control Hyperglycemia	Antidiabetic
Prevention and treatment of angina attacks	Antianginal
Prevention and treatment of thromboembolism and treatment of unstable angina and myocardial infarction	Antiplatelet
Reduction in frequency and severity of epileptic seizures	Anticonvulsant
Relief of moderate or severe pain	Opioid analgesic

2

Relief of extrapyramidal side effects	Antiparkinsonian
Treatment of primary hypercholesterolemia and mixed dyslipidemia	Lipopenic
Treatment of hypertension and edema due to heart failure and other causes	Diuretic
Treatment of hypertension, angina pectoris and heart failure	Beta blocker
Treatment of reversible airway obstruction due to asthma and COPD	Bronchodilator

3

Treatment of various forms of mood disorders	Antidepressant
Treatment of mild to moderate pain, fever and various inflammatory diseases	Non-steroidal anti-inflammatory drug (NSAID)
Treatment of nausea and vomiting	Antiemetic
Treatment of acute and chronic psychosis, and treatment of schizophrenia	Antipsychotic
Treatment of generalized anxiety disorder (GAD)	Anxiolytic

4

Diuretic	Hypokalemia
Beta-blocker	Bradycardia
Lipopenic	Abdominal distention
Opioid analgesic	CNS depression

5

Antihypertensive	Hypotension
Antiulcer	Constipation
Short-acting bronchodilator	Tremors/Shaking
Corticosteroid	Oral candidiasis

6

Antacid	Constipation
Antiplatelet	Gastrointestinal bleeding
Antianginal	Headache
Antiemetic	Drowsiness

7

Antidepressant	Weight gain
Antidiabetic	Tremors/Shaking
Anxiolytic	Anxiolytic
Laxative	Light cramps

8

Antibiotic	Administer in the conjunctival sac
Oral antidiabetic	Check for signs of hypoglycemia
Cholinesterase inhibitors	Regularly check BP & pulse
Anticoagulant	Checking for bruising

9

Analgesic	Assess pain before administering
Anticholinergic	Advise the patient to frequently rinse her mouth
Anxiolytic	Note the degree of sedation
Stool softener	Monitor abdominal distention

10

NSAID	Asking the beneficiary if her joint pain has decreased
Antiarrhythmic	Verifying the apical pulse before administration
Antidiabetic	Having a colleague verify the dose before administration
Antiulcer	Verifying whether the beneficiary's epigastric pain is still present

